



RECTOCELE

What is a rectocele?

A rectocele is a bulge of the front wall of the rectum into the vagina. The rectal wall may become thinned and weak, and it may balloon out into the vagina when you push down

to have a bowel movement. Most rectoceles occur in women where the front wall of the rectum is up against the back wall of the vagina. This area is called the rectovaginal septum and may be a weak area in the female anatomy. Other structures may also push into the vagina. The bladder bulging into the vagina is called a *cystocele*. The rectum bulging into the vagina is termed a *rectocele*. And the small intestines pushing down on the vagina from above may form an *enterocele*. Although uncommon, men may also develop a rectocele.

A rectocele may be present without any other abnormalities. In some cases, a rectocele may be part of a more generalized weakness of pelvic support

and may exist along with a cystocele, urethrocele, and enterocele, or with uterine or vaginal prolapse, rectal prolapse, and fecal or urinary incontinence.

What can cause a rectocele?

The underlying cause of a rectocele is a weakening of the pelvic support structures and thinning of the rectovaginal septum. Certain factors may increase the risk of a woman developing a rectocele. These include birth trauma such as multiple, difficult or prolonged deliveries, the use of forceps or other assisted methods of delivery, perineal tears, or an episiotomy into the rectum or anal sphincter muscles. In addition, a history of constipation and straining with bowel movements, or hysterectomy may contribute to the development of a rectocele. Commonly, these problems develop with age but they may occasionally occur in younger women or in those that have not delivered children.

What are the symptoms of a rectocele?

Many women have rectoceles but only a small percentage of women have symptoms related to the rectocele. Symptoms may be primarily vaginal or rectal. Vaginal symptoms include vaginal bulging, the sensation of a mass in the vagina, pain with intercourse or even something hanging out of the vagina that may become

irritated. Vaginal bleeding is occasionally seen if the vaginal lining of the rectocele is irritated, but other sources of the bleeding should be checked by your doctor. Rectal symptoms include constipation, particularly difficult evacuation with straining. Often this is associated with bulging in the vagina when straining to have a bowel movement. Some women find that pressing against the lower back wall of the vagina or along the rim of the vagina helps to empty the rectum. At times, there will be a rapid return of the urge to have a bowel movement after leaving the bathroom because stool that was trapped in the rectocele may return to the low rectum after standing up. A general feeling of pelvic pressure or discomfort is often present but this may be due to a variety of problems.

How is a rectocele diagnosed?

Most rectoceles may be identified on a routine office examination of the vagina and rectum. However, it may be difficult to assess the size and significance of the rectocele. A more accurate method of assessing the rectocele is an x-ray study called a defecagram. This study shows how large the rectocele is and if it empties with evacuation.

When should a rectocele be treated?

You should consider having your rectocele treated when it causes significant symptoms. It takes an experienced doctor to help you decide

whether your symptoms are caused by a rectocele. If there are multiple abnormalities present, it may be best to address them all at once as this will result in the best chance for improvement.

What treatment is available for a rectocele?

Rectoceles that are not causing symptoms do not need to be treated. In general, you should avoid constipation by eating a high fiber diet and drinking plenty of fluids.

Medical treatment

A bowel management program is the best first step. This includes a diet high in fiber and 6 to 8 glasses of fluids each day. Fiber acts like a sponge. It soaks up fluid so that less is removed as the stool travels around the colon. The stools will be larger, softer and easier to pass. You may wish to add a fiber supplement and/or a stool softener to this regimen to improve stool consistency. Most fiber supplements are made of psyllium, a seed product, or of a hydrophilic colloid (gel) that absorbs water. Most stool softeners are composed of docusate. This helps to smooth and lubricate the stool. Active laxatives are best avoided in most cases.

Avoid prolonged straining. If you cannot completely empty, get up and return later. Holding pressure with a finger to support the rectocele and

encourage the stool to go in the correct direction is often helpful. This may be accomplished by pressing against the lower back wall of the vagina or along the posterior rim of the vagina. Avoid placing a finger inside the anus to pull the stool out as this may cause harm. A pessary may be used to support the pelvic organs. It is a ring that is inserted into the vagina and must be individually fit to each woman.

Surgical treatment

If symptoms persist even with medical therapy, then surgical repair may be indicated. There are several surgical techniques used to repair a rectocele. A rectocele repair may be performed through the anus, through the vagina, through the perineum between the anus and vagina, or from above through the abdomen. When there is extensive pelvic relaxation and prolapse, the best approach may be a combined repair.

Who should treat me for this problem?

Both colorectal surgeons and gynecologists are trained to deal with these problems. If the symptoms are entirely vaginal, then it is appropriate for your gynecologist to address the problem. If your symptoms are rectal, then a colorectal surgeon should be involved. If there is any question, seek opinions from physicians of both specialties.

PATIENT INFORMATION

Rectocele



University Surgical Associates

Exceptional surgeons. Compassionate care.

Richard A. Moore, MD, FACS
J. Daniel Stanley, MD, FACS
Shauna Lorenzo-Rivero, MD, FACS
979 E. 3rd St., Suite C-300
Chattanooga, TN 37403
423.267.0466